COQ2 gene

coenzyme Q2, polyprenyltransferase

Normal Function

The *COQ2* gene provides instructions for making an enzyme called coenzyme Q2. This enzyme carries out one step in the production of a molecule called coenzyme Q10, which has several critical functions in cells throughout the body. In cell structures called mitochondria, coenzyme Q10 plays an essential role in a process called oxidative phosphorylation, which converts the energy from food into a form cells can use. Coenzyme Q10 is also involved in producing pyrimidines, which are building blocks of DNA, its chemical cousin RNA, and molecules such as ATP and GTP that serve as energy sources in the cell. In cell membranes, coenzyme Q10 acts as an antioxidant, protecting cells from damage caused by unstable oxygen-containing molecules (free radicals), which are byproducts of energy production.

Health Conditions Related to Genetic Changes

multiple system atrophy

Several variations in the *COQ2* gene have been suggested to increase the risk of multiple system atrophy, a progressive brain disorder that affects movement and balance and disrupts the function of the autonomic nervous system. The autonomic nervous system controls body functions that are mostly involuntary, such as regulation of blood pressure.

The identified variations alter single protein building blocks (amino acids) in coenzyme Q2. Most of the variations are very rare, but a genetic change that replaces the amino acid valine with the amino acid alanine at position 393 (written as Val393Ala or V393A) is relatively common. Studies suggest that these variations, including V393A, are associated with an increased risk of developing multiple system atrophy in the Japanese population. However, studies have not found a correlation between COQ2 gene variations and multiple system atrophy in other populations, including Koreans, Europeans, and North Americans. It remains unclear whether COQ2 gene variations represent a significant risk factor for this disease.

Researchers speculate that changes in the *COQ2* gene could impair the activity of coenzyme Q2, which would affect the production of coenzyme Q10. Levels of coenzyme Q10 are reduced in the brains of people with multiple system atrophy. A reduction in the amount of coenzyme Q10 may impair cellular energy production from oxidative phosphorylation and increase the vulnerability of cells to damage from

free radicals. However, it is unknown how these changes are related to the specific features of this disorder.


other disorders

At least six mutations in the *COQ2* gene have been found to cause a disorder known as primary coenzyme Q10 deficiency. This rare disease, which appears in infancy or early childhood, can affect many parts of the body, most often the brain, muscles, and kidneys. The *COQ2* gene mutations associated with this disorder greatly reduce or eliminate the production of coenzyme Q2, which prevents the normal production of coenzyme Q10. These mutations are thought to affect coenzyme Q10 levels more severely than the *COQ2* gene variations associated with multiple system atrophy (described above). Studies suggest that a shortage (deficiency) of coenzyme Q10 impairs oxidative phosphorylation and increases the vulnerability of cells to damage from free radicals. A deficiency of coenzyme Q10 may also disrupt the production of pyrimidines. These changes can cause cells throughout the body to malfunction, which may help explain the variety of organs and tissues that can be affected by primary coenzyme Q10 deficiency.

Chromosomal Location

Cytogenetic Location: 4q21.22-q21.23, which is the long (q) arm of chromosome 4 between positions 21.22 and 21.23

Molecular Location: base pairs 83,138,568 to 83,285,129 on chromosome 4 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- 4-HB polyprenyltransferase
- 4-hydroxybenzoate decaprenyltransferase
- 4-hydroxybenzoate polyprenyltransferase, mitochondrial
- CL640
- coenzyme Q2 4-hydroxybenzoate polyprenyltransferase

- coenzyme Q2 homolog, prenyltransferase
- COQ10D1
- FLJ26072
- MSA1
- para-hydroxybenzoate-polyprenyltransferase, mitochondrial
- PHB:polyprenyltransferase
- PHB:PPT

Additional Information & Resources

Educational Resources

- Linus Pauling Institute, Oregon State University: Coenzyme Q10 http://lpi.oregonstate.edu/mic/dietary-factors/coenzyme-Q10
- Molecular Biology of the Cell (fourth edition, 2002): How Cells Obtain Energy from Food
 - https://www.ncbi.nlm.nih.gov/books/NBK26882/
- The Cell: A Molecular Approach (second edition, 2000): The Mechanism of Oxidative Phosphorylation https://www.ncbi.nlm.nih.gov/books/NBK9885/

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28COQ2%5BTIAB%5D%29+OR+%28coenzyme+Q2%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D

OMIM

 COQ2, S. CEREVISIAE, HOMOLOG OF http://omim.org/entry/609825

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology http://atlasgeneticsoncology.org/Genes/GC_COQ2.html
- ClinVar https://www.ncbi.nlm.nih.gov/clinvar?term=COQ2%5Bgene%5D
- HGNC Gene Symbol Report http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/ hgnc_data.php&hgnc_id=25223

- NCBI Gene https://www.ncbi.nlm.nih.gov/gene/27235
- UniProt http://www.uniprot.org/uniprot/Q96H96

Sources for This Summary

- OMIM: COQ2, S. CEREVISIAE, HOMOLOG OF http://omim.org/entry/609825
- Mitsui J, Tsuji S. Mutant COQ2 in multiple-system atrophy. N Engl J Med. 2014 Jul 3;371(1):82-3. doi: 10.1056/NEJMc1311763.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/24988566
- Mollet J, Giurgea I, Schlemmer D, Dallner G, Chretien D, Delahodde A, Bacq D, de Lonlay P, Munnich A, Rötig A. Prenyldiphosphate synthase, subunit 1 (PDSS1) and OH-benzoate polyprenyltransferase (COQ2) mutations in ubiquinone deficiency and oxidative phosphorylation disorders. J Clin Invest. 2007 Mar;117(3):765-72.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/17332895
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1804361/
- Multiple-System Atrophy Research Collaboration. Mutations in COQ2 in familial and sporadic multiple-system atrophy. N Engl J Med. 2013 Jul 18;369(3):233-44. doi: 10.1056/NEJMoa1212115. Epub 2013 Jun 12. Erratum in: N Engl J Med. 2014 Jul 3;371(1):94. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/23758206
- Schottlaender LV, Bettencourt C, Kiely AP, Chalasani A, Neergheen V, Holton JL, Hargreaves I, Houlden H. Coenzyme Q10 Levels Are Decreased in the Cerebellum of Multiple-System Atrophy Patients. PLoS One. 2016 Feb 19;11(2):e0149557. doi: 10.1371/journal.pone.0149557. eCollection 2016.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/26894433
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4760984/
- Sun Z, Ohta Y, Yamashita T, Sato K, Takemoto M, Hishikawa N, Abe K. New susceptible variant of COQ2 gene in Japanese patients with sporadic multiple system atrophy. Neurol Genet. 2016 Mar 3; 2(2):e54. doi: 10.1212/NXG.000000000000054. eCollection 2016 Apr. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/27123473
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4830192/
- Zhao Q, Yang X, Tian S, An R, Zheng J, Xu Y. Association of the COQ2 V393A variant with risk of multiple system atrophy in East Asians: a case-control study and meta-analysis of the literature. Neurol Sci. 2016 Mar;37(3):423-30. doi: 10.1007/s10072-015-2414-8. Epub 2015 Nov 21. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/26590992

Reprinted from Genetics Home Reference: https://ghr.nlm.nih.gov/gene/COQ2 Reviewed: July 2016

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services